21-283-9040

Oct 10, 2021
RECEIVED DATE
Oct 08, 2021

SEND TO **35432** 



PAGE 1/2

Oct 10, 2021

13611 B Street • Omaha, Nebraska 68144-3693 • (402) 334-7770 www.midwestlabs.com

# REPORT OF ANALYSIS

For: (35432) OPEN FARM Dog-OF Turkey & Chicken Kibble 27821FTC

Analysis	Level Found As Received	Units	Reporting Limit	Method	Analyst- Date	Verified- Date
Sample ID: 27821FTC - Micros	Lab Number: <b>13706598</b> D	oate Sampled: 20	21-10-05			
Salmonella	negative	org/375g	1	RapidChek/AOAC RI 030301; AFNOR SDI 34/01-04/10	kkb0-2021/10/10	jzh4-2021/10/10
E. coli (generic)	n.d.	cfu/g	10	AOAC 2018.13	kje1-2021/10/09	jzh4-2021/10/09
Listeria	negative	org/125g	1	RapidChek/AOAC RI 020401	kkb0-2021/10/10	jzh4-2021/10/10

All results are reported on an AS RECEIVED basis., n.d. = not detected, cfu = colony forming unit

For questions please contact:

Derrick Kendrick Account Manager

dkendrick@midwestlabs.com (402)590-2989

21-283-9040

Oct 10, 2021 RECEIVED DATE Oct 08, 2021

35432



**PAGE 2/2** Oct 10, 2021

13611 B Street • Omaha, Nebraska 68144-3693 • (402) 334-7770 www.midwestlabs.com

## REPORT OF ANALYSIS

For: (35432) OPEN FARM - Dog-OF Turkey & Chicken Kibble 27821FTC

# **Detailed Method Description(s)**

## E. coli and Total Coliform using 3M Petrifilm

Sample analysis follows MWL MI 292 which is based on AOAC 2018.13. A representative sample is obtained and added to phosphate buffer. Aliquots of the sample are withdrawn and placed on Petrifilm plates. The plates are incubated for 18 to 24 hours. After incubation, the plates are counted to determine the number of generic E. coli and total coliforms present. The color of the colony and the presence of gas differentiate a generic coliform from E. coli. The levels are reported as colony forming units (cfu).

#### Salmonella - Lateral Flow

Samples are analyzed following MWL MI 195 which is based on the RapidChek Select Salmonella User Guide. A representative sample is obtained and combined with a selective media and allowed to incubate. After incubation, a test strip is used for Salmonella determination. Results are reported as negative or presumptive positive.

### Listeria Lateral Flow

Samples are analyzed following MWL MI 194 which is based on the RapidChek Listeria User Guide. A representative sample is obtained and combined with a selective growth media. It is incubated for 40-48 hours. After incubation, an aliquot is heated for 10 minutes, and a test strip for Listeria detection is used. Results are reported as negative or presumptive positive. This procedure does not speciate Listeria.